



FACULDADE DE MEDICINA
UNIVERSIDADE DO PORTO

MESTRADO INTEGRADO EM MEDICINA

2011/2012

Joana Vicente Vargas

Evaluation of Central Nervous System involvement in SLE patients.

Screening psychiatric manifestations – A Systematic Review

março, 2012

FMUP



FACULDADE DE MEDICINA
UNIVERSIDADE DO PORTO

Joana Vicente Vargas

Evaluation of Central Nervous System involvement in SLE patients.

Screening psychiatric manifestations – A Systematic Review

Mestrado Integrado em Medicina

Área: Reumatologia

Trabalho efetuado sob a Orientação de:

Doutor Carlos Jorge Cabral Vaz

Trabalho organizado de acordo com as normas da revista:

Acta Reumatológica Portuguesa

março, 2012

FMUP

Eu, Joana Vicete Vargas abaixo assinado, nº mecanográfico 060801038, estudante do 6º ano do Mestrado Integrado em Medicina, na Faculdade de Medicina da Universidade do Porto, declaro ter atuado com absoluta integridade na elaboração deste projeto de opção.

Neste sentido, confirmo que **NÃO** incorri em plágio (ato pelo qual um indivíduo, mesmo por omissão, assume a autoria de um determinado trabalho intelectual, ou partes dele). Mais declaro que todas as frases que retirei de trabalhos anteriores pertencentes a outros autores, foram referenciadas, ou redigidas com novas palavras, tendo colocado, neste caso, a citação da fonte bibliográfica.

Faculdade de Medicina da Universidade do Porto, 16/03/2012

Assinatura:

Joana Vicete Vargas

Nome: Joana Vicente Vargas

Endereço eletrónico: joanavargas@gmail.com

Telefone ou Telemóvel: 918778831

Número do Bilhete de Identidade: 13435784

Monografia:

Evaluation of Central Nervous System involvement in SLE patients. Screening psychiatric manifestations – A Systematic Review

Orientador:

Doutor Carlos Jorge Cabral Vaz

Ano de conclusão: 2012

Designação da área do projeto:

Reumatologia

É autorizada a reprodução integral desta Dissertação/Monografia (cortar o que não interessar) para efeitos de investigação e de divulgação pedagógica, em programas e projetos coordenados pela FMUP.

Faculdade de Medicina da Universidade do Porto, 16/03/2012

Assinatura:

Joana Vicente Vargas

*Agradeço ao Doutor Carlos Vaz pela
ajuda e orientação dada ao longo deste trabalho*

Evaluation of Central Nervous System involvement in SLE patients.

Screening psychiatric manifestations – A Systematic Review

Avaliação do envolvimento do Sistema Nervoso Central em doentes lúpicos.

Rastreio de manifestações psiquiátricas – Revisão Sistemática

Joana Vicente Vargas

Aluna do 6º ano do curso de Mestrado Integrado de Medicina
da Faculdade de Medicina da Universidade do Porto

Serviço de Reumatologia do Hospital de São João
Faculdade de Medicina da Universidade do Porto

Alameda Prof. Hernâni Monteiro
4200-319 Porto, Portugal

med06038@med.up.pt

Evaluation of Central Nervous System involvement in SLE patients. Screening psychiatric manifestations – A Systematic Review

Avaliação do envolvimento do Sistema Nervoso Central em doentes lúpicos. Rastreamento de manifestações psiquiátricas – Revisão Sistemática

Abstract: Cognitive dysfunction, mood and anxiety disorders are three out of the five psychiatric syndromes included in Neuropsychiatric Lupus. These manifestations are among the most prevalent in SLE having an important impact on patients quality of life. However, the unknown etiology allied to the lack of clarity on the best diagnosis procedure, makes early diagnosis difficult. This manuscript reviews the recent literature on the screening instruments focused on identifying lupus patients with probable psychiatric manifestations.

Resumo: Disfunção cognitiva, distúrbios do humor e ansiedade são três das cinco síndromes psiquiátricas incluídas na Lupus Neuropsiquiátrica. Estas manifestações estão entre as mais prevalentes na lúpus e têm um impacto importante na qualidade de vida destes doentes. Apesar disto, o desconhecimento etiológico aliado à falta de um processo diagnóstico ideal, fazem com que o seu diagnóstico seja por vezes difícil. Neste trabalho é apresentada uma revisão da literatura recente dos instrumentos de rastreio da identificação de doentes com lúpus com prováveis manifestações psiquiátricas.

Key-words: Neuropsychiatric Lupus, Screening, Cognitive Dysfunction, Mood Disorders, Anxiety Disorders.

Palavras-Chave: Lupus Neuropsiquiátrica, Rastreamento, Disfunção Cognitiva, Distúrbios do Humor, Distúrbios da Ansiedade.

INTRODUCTION

Systemic Lupus Erythematosus (SLE) is a chronic, inflammatory, autoimmune disease characterized by its multi-systemic involvement and multiple clinical manifestations. It can involve the nervous system on its central or peripheral component.

In an effort to standardize nomenclature and diagnostic methodology, the American College of Rheumatology Ad Hoc Committee on Neuropsychiatric Lupus, published in 1999, the case definitions and diagnostic recommendations on Neuropsychiatric Lupus (NPSLE). Nineteen syndromes were included in NPSLE and for each were suggested not only the diagnostic criteria and testing but also associations and exclusions¹. The publication was widely accepted and has been used ever since in the study of NPSLE in adult and pediatric populations.

Recent studies report a prevalence of neuropsychiatric manifestations of 27-80% in adults²⁻⁸, and 22-95% in children⁹⁻¹². NPSLE can develop in any time during the course of the disease but there are several studies reporting a tendency to occur early in its course^{5,8}. Furthermore, association between neuropsychiatric events and disease activity has been reported in some studies^{13, 14} and denied in others¹⁵⁻¹⁹.

Five out of 19 syndromes that included in NPSLE are psychiatric and were considered by the ACR Ad Hoc Committee on NPSLE according to the DSM-IV diagnostic criteria: mood disorders, anxiety disorders, cognitive dysfunction, psychosis and acute confusional state.

The importance of studying psychiatric phenomena in lupus disease lies not only in its high prevalence reported²⁰ but mainly in what it represent clinically and socioeconomically. Psychiatric manifestations have been associated with a decreased

quality of life^{5,8,19}, increased functional disability³, sleep disorders^{21,22}, increased unemployment rate^{23,34} and health service utilization^{25,26}.

There is still little consensus on the role of laboratory tests and imaging techniques in the diagnosis of psychiatric syndromes and due to the lack of simple diagnosis process, psychiatric syndromes are frequently undiagnosed.

This manuscript pretends to be a systematic review on the recent literature focused in the study of screening tools for the identification of SLE patients with probable cognitive dysfunction, mood disorders or anxiety disorders, the three most prevalent psychiatric manifestations in lupus.

METHODS

A Pubmed/Medline and Scopus search was conducted from January 2002 to January 2012 using the following keywords: “neuropsychiatric lupus” or “systemic lupus erythematosus” combined with “diagnosis”, “anxiety disorders”, “mood disorders” and “cognitive dysfunction”. A follow-up of the relevant bibliography in articles was also done in order to identify additional relevant studies.

Abstracts of all identified studies were reviewed by two investigators. Every time an abstract was considered as potentially relevant, by either or both investigators, the full-text was retrieved and reviewed for relevance by applying the exclusion and inclusion criteria.

Review articles, case reports, studies in languages other than english or portuguese were immediately excluded. Included were all the articles investigating a screening tool for the diagnosis of psychiatric lupus. Studies that did not relate with diagnosis of psychiatric

manifestations in the lupus setting or with its laboratory and/or imaging diagnosis were also excluded.

RESULTS

The described search identified 446 articles (Figure I), of which 102 were considered as potentially relevant. Forty-seven articles were excluded based on abstract analysis. The remaining 55 studies were reviewed on its full text and the inclusion and exclusion criteria were applied. Thirteen studies were included in this systematic review.

COGNITIVE DYSFUNCTION

Difficulties in remembering, concentrating and performing cognitive-dependent activities are frequent complaints of SLE patients^{27,28}. In fact Cognitive Dysfunction (CD) has been reported as one of the most frequent neuropsychiatric manifestations in SLE, having a prevalence in the adult population of 5,4 - 50 %^{5,6,14,16,28-33} and in the pediatric population of 7,3 - 79,8 %^{5,6,14,19,29,30}.

In adults CD increases the risk of physical injury, reduces patients ability to properly adhere to treatment regimens and to function effectively in their home or work environments¹⁷. In pediatric patients cognitive impairment may prevent the normal development, with serious repercussions throughout life³⁴.

A comprehensive battery of neuropsychological tests is the ideal method to evaluate the presence and severity of cognitive dysfunction. However, in order to facilitate the diagnosis process, the ACR Ad Hoc Committee on NPSLE proposes an one-hour battery of brief mental status examinations (short ACR-SLE battery)¹.

The short ACR-SLE battery has been validated and found reliable in spite of high practice-effect observed in some tests by a study that tested the correlation between the short ACR-SLE battery and a comprehensive neuropsychological battery³⁵. Nevertheless the short ACR-SLE battery is not easily available, requires administration by specialized professionals and is too expensive to be used in routine clinical consults^{15,27,28}.

The ANAM is a, 30 to 45 minutes self-administered, computerized battery of neuropsychological tests, developed by U.S. military in order to assess the cognitive repercussions of chemical agents, extreme environments and fatigue on cognitive processing speed and efficiency²⁷.

Holliday et al, in 2003, was the first to suggest the use of ANAM as a screening tool in the SLE context. This study administered both ANAM and a traditional test battery based on ACR Ad Hoc Committee on NPSLE recommendations in a sample of 67 ethnically mixed SLE patients enrolled in a large prospective cohort²⁷.

The results showed that many ANAM measures correlated with the scores from the traditional neuropsychological tests. It was also found that age is of little relevance on the variance observed when accounted alone but, it acts as a powerful moderator variable when is entered with ANAM variables into a linear regression model. This model accounted for about 61% of the variance in the average T-score on the traditional tests²⁷.

A Roebuck-Spencer, 2006 publication, confirms the positive correlation between ANAM and traditional neuropsychological testing. In this study the performances in the two batteries are compared in a sample of 60 SLE patients participating in a large SLE cohort on biomarkers of cognitive dysfunction. ANAM test battery demonstrated a sensibility of 76,2% and specificity of 82,8% on the classification of individuals with probable cognitive impairment versus no impairment in neuropsychological testing. ANAM remained a good

screening tool when depression and sleepiness were present, measured with validated self-reported measures of sleepiness and depressed mood, suggesting that it is not confounded by these. Furthermore, significant correlation between ANAM's mood scale and BDI-II was found, supporting its use as a potential measure of emotional distress in SLE patients³⁶.

A 2010 study by Hanly et al, sought out to compare ANAM battery tests' performance in a sample of 29 healthy controls, 68 Lupus (SLE), 33 Rheumatoid Arthritis (RA) and 20 Multiple Sclerosis (MS) patients³¹.

The results showed a cognitive impairment of 11-50% in SLE patients (depending on stringency of classification rules) when compared with locally recruited healthy controls. However this frequency was comparable with the 9-61% calculated frequency in RA patients and lower than that calculated for MS patients, 20-75%. The frequency difference between SLE patients and patients with stable MS disease is expected but the observed comparability of frequencies between SLE patients and patients with a disease that does not affect primarily the CNS, as RA, raised questions about the presumed etiology of deficits detected by ANAM. The authors suggested that the measures evaluated by the ANAM battery do not distinguish between impaired mental processing and speed sensorimotor deficiencies and can instead represent CNS immunosuppressive toxicity. Leading to the conclusion that ANAM can't be used to determine dysfunction on specific cognitive domains and it was not designed as a substitute for formal neuropsychological assessment³¹.

In 2007 Brunner et al studied the statistical properties of the pediatric ANAM (ped-ANAM) in a childhood-onset SLE sample. Ped-ANAM and a battery of formal neuropsychological tests (based on published data for SLE adults) were performed in a

sample of 27 children with a median age of 16,5 years recruited from a pediatric rheumatology clinic. A trend towards worse performance of participants with CD compared to those without was observed in every performance parameter of the battery but statistically significant differences between the two groups were only reached for 3 out of the 10 ped-ANAM tests. Furthermore, statistically significant correlations were found between Ped-ANAM tests and formal neuropsychological tests. Ped-ANAM was found as a promising tool to screen cognitive dysfunction in SLE children presenting validity and promising sensitivity and specificity¹⁵.

The Cognitive Symptom Inventory (CSI) is a self-administered paper questionnaire consisting of 21 items focused in evaluating the subject's ability to perform several cognitive functions and activities of daily life.

In 2002, Alarcón et al published a study which aimed determine the factor structure of the CSI and the production of 4 factor scales and their correlation with 3 self-report measures of cognitive dysfunction and self-report measures of fatigue, helplessness, self-efficacy, pain, social support and use of maladaptive coping skills. The sample, drawn from a large prospective cohort (LUMINA), consisted in 156 ethnically mixed SLE patients¹⁷.

The four main factors assessed by the CSI were found to be: Attention/Concentration, Pattern/Activity Management, Intermediate Memory and Initiation of Executive Functions. Despite the small shared amount of common variance between these four factors, the correlation was not high enough that they duplicate one another¹⁷.

Modest statistically significant correlations were also found between CSI cognitive factor scales and SLAM measure of Cortical Dysfunction, SF-36 measure of Mental Functioning,

SDI measure of Cognitive Impairment, measures of fatigue, psychological distress, social support, maladaptive coping skills, self-efficacy and pain¹⁷.

CSI patients' responses were found not to be confounded by social-demographic or clinical variables. The questionnaire was completed in an average time of 10 minutes and with minimal paraprofessional help regardless of ethnical backgrounds or administered language¹⁷.

The Montreal Cognitive Assessment (MoCA) is a validated, one-page, physician-administered questionnaire used on the identification of mild cognitive dysfunction in the elderly³².

Published in 2011 by Adhikari et al there is a study that aims to evaluate MoCA as a screening tool for detection of cognitive dysfunction in SLE patients. In a sample of 44 SLE patients recruited in the Cincinnati area (USA) were applied both the MoCA and, as gold standard, the ANAM³⁵. Results demonstrate that to a standard cutoff score of 26 the sensitivity of MoCA was 83%. The specificity was 73% with a positive predictive value of 50% and a negative predictive value of 92%. These results suggest that MoCA has the potential to be used as a screening tool for the detection of SLE with probable cognitive dysfunction³².

In 2008, Kozora et al publishes a study aimed to examine the screening utility of the standardized neurologic evaluations in the identification of SLE patients with probable cognitive dysfunction. All the participants in the study were already enrolled in a large prospective cohort of cognitive functioning and neuroimaging. The participants were selected based on the examination of their clinical history and on physician interview in order to identify the ones with history of neuropsychiatric diseases or depression. The Scripps Neurologic Rating Scale (NRS) and the short ACR-SLE battery were administered

to all the participants (SLE=67, Controls=29)³⁷. The NRS is a 22 item neurologic exam developed for the clinical evaluation of patients with multiple sclerosis.

The prevalence of cognitive dysfunction in the sample was 20,9%. The non-NPSLE group had worse outcomes on NRS global score than the control group ($p<0,001$). However after analysis of the NRS parameters, the one responsible for the statistically significant difference was “mentation and mood”. Nevertheless two patients were excluded after the initial screening process during the administration of the neurologic examination by the neurologist suggesting that the NRS can assure that overt neurologic dysfunction is not present and assist in identifying non-NPSLE patients³⁷.

Julian et al publishes, in 2011, a study aimed at the evaluation of the utility of telephone screening and self-report assessments of cognitive complaints in detecting cognitive impairment in individuals with SLE and RA²⁸.

Two screening measures were evaluated: a 12-15 minutes telephone interview based on three neuropsychological tests (see article for details) and the Perceived Deficits Questionnaire (PDQ), a five-question, self-administered questionnaire. A validated neuropsychological battery based on the short ACR-SLE battery was used as “gold standard”.

The sample of 138 SLE patients and 84 RA patients was drawn from two large cohorts of SLE and RA patients, respectively. The cognitive dysfunction rates were 27% to the SLE group. The telephone screening had 77% sensitivity, 65% specificity, 94% negative predictive value, 43% positive predictive value and 67% of the patients were correctly classified in the SLE group. While the PDQ had 64% sensitivity, 65% specificity, 83% negative predictive value, 38% positive predictive value and 64% of the patients were correctly classified in the SLE group.

Contrary to telephone screening measure, the PDQ was not a significant predictor of cognitive impairment when adjusted for social-demographic data and depression²⁸.

MOOD AND ANXIETY DISORDERS

The reported prevalence of mood and anxiety disorders in recent studies ranges between 12,4-60%^{5,14,28,38-41} and 6,4-46,5%^{5,29,39,41}, respectively.

The ACR Ad Hoc Committee on NPSLE recommends the use of standardized instruments like the Center for Epidemiological Studies - Depression Scale (CES-D) and the Hospital Anxiety and Depression Scale (HADS) for the diagnosis of mood and anxiety disorders¹.

Several associations have been sought out by different studies in order to understand the pathophysiology of these disorders.

The association between short disease duration and anxiety disorders was described in a 2011 study by Hawro et al, raising the hypothesis that anxiety was a consequence of the inadequate information about the disease suggesting that at least part of the anxious disorders encountered in NPSLE have an adaptive background³³.

Kozora et al in 2007 compared the performances of depressed SLE patients (n=13), depressive non-SLE patients (n=10) and healthy controls (n=25) in the short ACR-SLE battery and a comprehensive neuropsychological battery. The results of this study not only confirmed the association between depression and cognitive dysfunction⁴² but also validated the short ACR-SLE battery for the diagnosis of cognitive dysfunction in depressed SLE patients⁴³.

In a survey study published by Iverson et al in 2002, two screening depression measures were compared, Beck Depression Inventory-Second Edition (BDI-II) and the British Columbia Major Depression Inventory (BCMDI), both instruments constructed according

to the diagnostic criteria of DSM-IV for depression. The sample consisted of 103 self-reported lupus patients (no attempt was made to confirm the diagnosis) and 136 healthy controls. The self-reported SLE group had higher rates of depression and vegetative symptoms (fatigue, difficulty falling asleep, sadness, etc.) than the control group. The results point to a possible over-estimate diagnosis of depression in this study of 103 SLE patients, as measured by the BDI-II, a test that is considered valid and reliable screening of depression. Fifteen percent of the patients identified as depressed on the BDI-II scored in the normal range on the BCMDI, and 46% scored in the possibly depressed range. Therefore, it is possible that the BDI-II over-identified depression in this sample²².

In a 2011 study by Julian et al, the Center for Epidemiological Studies - Depression Scale (CES-D) is compared with the Mini-International Neuropsychiatric Interview (MINI), a validated diagnostic method based on structured clinical interview, in a sample of 150 SLE patients drawn from a prospective cohort of 957 lupus patients. In this sample 26% of the patients were diagnosed with a mood disorder and 17% with major depressive disorder measured by the MINI. The results showed a 92% of corrected classified patients with major depressive disorder and a 87% sensibility and specificity in detection of any mood disorder with the CES-D⁴⁰.

Hyphantis et al, 2011, studies the psychometric characteristics of the greek version of the Patient Health Questionnaire-9 (PHQ-9) in a diverse rheumatologic sample of 558 patients (62 lupus patients). The PHQ-9 is a, 9 question, self-administered, simple, questionnaire used on the screening and severity assessment of depression. The results showed 25,4% prevalence of Major Depressive Disorder, measured by MINI, and a 81,2% and 86,8%, sensitivity and specificity, respectively, of the greek version PHQ-9⁴⁴.

PILOT SCREENING TOOL

Mosca et al published, in 2011, a study aimed to create a questionnaire, to be administered by physicians, that could be used as a screening tool for the identification of neuropsychiatric manifestations in SLE patients with no obvious CNS involvement, for further evaluation⁴⁵.

Starting from group of 112 questions drawn from 41 questionnaires aimed at assessing neuropsychiatric manifestations similar to those prevalent in NPSLE, a panel of experts and statistic analysis created a draft questionnaire with 62 items. This draft questionnaire was then tested in 139 SLE patients from 11 european centers and the results were compared with clinical diagnosis made by a specialist. After additional statistical analysis, the final questionnaire consisted of 27 items, 15 referring to the central nervous system symptoms and 12 to psychiatric ones.

For a cutoff value of 17 the questionnaire had a sensitivity and specificity of 93% and 25% respectively. It was found that the association of non-specific symptoms (e.g. subjective complain of cognitive dysfunction or headache) may determine a higher score than serious manifestations such as seizures alone. All the initial included items aimed at the assessment of peripheral nervous system symptoms were excluded based on the methodology used. Despite the intent to be administered by a physician the authors considered the questionnaire to be simple enough to be filled by the patient himself⁴⁵.

DISCUSSION

One of the most controversial topics in the study of the psychiatric manifestations is their etiology. In fact, despite the high prevalences reported its still not known if they are a direct consequence of the autoimmune disease or secondary to it. Several factors can

explain the secondary nature of these disorders like the stress of having a chronic disease, the lack of social support or the use of immunosuppressive therapy³⁹.

Psychiatric syndromes are rarely diagnosed early in their course due to their initially faint clinical manifestations and lack of accepted, valid and accessible methods of detection which leads to under-diagnosis and under-treatment of these conditions. Thus, a simple, sensitive screening test would serve to improve quality of management of lupus patients³². On the other hand, randomized clinical trials are necessary in SLE to fully understand the benefit-harm tradeoffs for screening these psychiatric manifestations⁴⁰.

Some instruments with the intend of screening cognitive dysfunction in SLE patients have been recently analyzed. However a lot more research needs to be done in order to validate them. With the exception of ANAM, all the instruments proposed as screening tests across the literature were studied only once.

The ANAM presents as the most analyzed of the screening tools, presenting good sensitivity and specificity (76,2% and 82,4%, respectively) on the distinction of SLE patients with cognitive dysfunction in formal neuropsychological testing from those without. ANAM presents as a instrument with less sensitivity to confounding variables such as, education, English proficiency and possible ethnic differences compared to the formal neuropsychological testing²⁷. Its accessibility, self-administration, low practice-effects, reduced cost and validity in depressed lupus patients^{27,36} makes of ANAM a promising screening. The finding that it may be confounded by immunosuppressive toxicity and/or sensorimotor deficits³¹ may not be as relevant clinically as it is etiologically since regardless of cause attribution, cognitive dysfunction is a co-morbidity that needs to be addressed when diagnosed.

Preliminary results determined that Cognitive Symptom Inventory (CSI), a 10 minutes self-administered paper test, has the potential to be used as a screening instrument on the identification of SLE patients with probable cognitive dysfunction. Revision of some of the items' content, expansion of the questionnaire so it covers other cognitive domains, study of its psychometric characteristics and testing it in different samples are some of the subjects that need to be addressed in further studies in order to validate the CSI as a screening tool¹⁷. Contrary to CSI, the Perceived Deficits Questionnaire, a 5 question self-administered questionnaire was found to be a weak predictor of cognitive impairment since it was confounded by social-demographic data and depression²⁸.

The Montreal Cognitive Assessment (MoCA), a physician administered questionnaire, presented a 83% sensitivity and 73% specificity. However the "gold standard" used was not formal neuropsychological testing but ANAM, which represents a vulnerability of the study³².

In the Kozora et al study on the utility as a screening tool of the Scripps Neurologic Rating Scale¹⁶ the results were not too impressive since despite the significant difference between SLE and control groups in identifying cognitive dysfunction, "mentation and mood" was the parameter responsible for the statistically significant difference. Furthermore, one aspect that was not referred in the study was how long did it take to administer the neurologic exam, essential to determine the true time-cost efficiency of it. Another promising screening instrument is the Telephone Screening studied by Julian et al, the 43% positive predictive value and 93% negative predictive value presented by this tool, is an advantage in that permits to exclude with greater confidence individuals without cognitive impairment²⁸.

It is somehow surprising that despite the high frequencies reported of mood and anxiety disorders among lupus patients, and the number of measures available aiming the screening of these disturbances, only two studies were found comparing the performances of these tests in the lupus context. In fact, only in 2011 the Center for Epidemiological Studies Depression Scale (CES-D), recommended in 1999 by the ACR Ad Hoc Committee on NPSLE as a screening tool to identify patients with probable mood disorders, was tested, presenting a 87% sensitivity and specificity in detecting any mood disorder when compared with MINI⁴⁰.

Studies like the one published by Hyphantis et al, that validated the greek version of PHQ-9 in a large and diverse rheumatologic sample⁴⁴, are very important since questionnaires when translated need to be validated again in order to guarantee their quality.

Pediatric Neuropsychiatric Lupus (Ped-NPSLE) is even less understood than the adult form due to the lack of studies published. The definition cases and diagnostic criteria and testing recommended by the ACR Ad Hoc Committee on NPSLE are being inadequately used in children. Williams et al demonstrated that depending on the methodology used to classify cognitive impairment in children, the prevalence of cognitive dysfunction ranged from 7,3% to 63,4% in the same sample. More than that no significant differences were encountered in tested domain scores as well as cognitive dysfunction prevalence estimates between lupus children and controls¹⁹. Another difficulty in studying pediatric lupus is the small sample available, one of the main limitations of the Brunner et al study¹⁵, that examined the usefulness of the pediatric version of ANAM in small sample of 27 lupus children.

Mosca et al approached screening testing in a holistic way, creating a physician-administered simple questionnaire to screen neuropsychiatric events. The main limitation of the study is the lack of items that assess the peripheral nervous system, justified by the authors as a result of the low prevalence of these phenomena in the lupus context. Despite the low specificity demonstrated (25%) this is a revolutionary study that demonstrated promising preliminary results⁴⁵.

There is still a long way to go regarding Neuropsychiatric Lupus. Sample selection, classification of impairment and attribution of cause are parameters that need to be defined and standardized in order to compare studies and draw conclusions.

The screening of psychiatric manifestations needs to be evaluated through randomized clinical trials in its true utility since the impact of early intervention on these phenomena is still unknown.

The translation and validation of screening tools in portuguese might facilitate and stimulate the study of neuropsychiatric lupus in Portugal.

REFERÊNCIAS

- 1- The American College of Rheumatology nomenclature and case definitions for neuropsychiatric lupus syndromes. *Arthritis Rheum* 1999; 42:599-608.
- 2 - Brey RL, Holliday SL, Saklad AR. Neuropsychiatric syndromes in lupus: prevalence using standardized definitions. *Neurology* 2002; 58:1214-1220.
- 3 - Jönsen A, Bengtsson AA, Nived O, Ryberg B, Sturfelt G. Outcome of neuropsychiatric systemic lupus erythematosus within a defined Swedish population: increased morbidity but low mortality. *Rheumatology (Oxford)* 2002; 41:1308-1312.

- 4 - Mikdashi J, Handwerger B. Predictors of neuropsychiatric damage in systemic lupus erythematosus: data from the Maryland lupus cohort. *Rheumatology (Oxford)* 2004; 43:1555-1560.
- 5 - Hanly JG, Urowitz MB, Sanchez-Guerrero J. Systemic Lupus International Collaborating Clinics. Neuropsychiatric events at the time of diagnosis of systemic lupus erythematosus: an international inception cohort study. *Arthritis Rheum* 2007; 56:265-273.
- 6 - Monov S, Monova D. Classification criteria for neuropsychiatric systemic lupus erythematosus: do they need a discussion? *Hippokratia* 2008; 12:103-107.
- 7 - Zhu TY, Tam LS, Lee VW, Lee KK, Li EK. Systemic lupus erythematosus with neuropsychiatric manifestation incurs high disease costs: a cost-of-illness study in Hong Kong. *Rheumatology (Oxford)* 2009; 48:564-568.
- 8 - Hanly JG, Urowitz MB, Su L; Systemic Lupus International Collaborating Clinics (SLICC). Prospective analysis of neuropsychiatric events in an international disease inception cohort of patients with systemic lupus erythematosus. *Ann Rheum Dis* 2010; 69:529-535.
- 9 - Sibbitt WL Jr, Brandt JR, Johnson CR. The incidence and prevalence of neuropsychiatric syndromes in pediatric onset systemic lupus erythematosus. *J Rheumatol.* 2002; 29:1536-1542.
- 10 - Olfat MO, Al-Mayouf SM, Muzaffer MA. Pattern of neuropsychiatric manifestations and outcome in juvenile systemic lupus erythematosus. *Clin Rheumatol* 2004; 23:395-399.
- 11 - Yu HH, Lee JH, Wang LC, Yang YH, Chiang BL. Neuropsychiatric manifestations in pediatric systemic lupus erythematosus: a 20-year study. *Lupus* 2006; 15:651-657. Review. Erratum in: *Lupus* 2007; 16:232.

- 12 - Hiraki LT, Benseler SM, Tyrrell PN, Hebert D, Harvey E, Silverman ED. Clinical and laboratory characteristics and long-term outcome of pediatric systemic lupus erythematosus: a longitudinal study. *J Pediatr* 2008; 152:550-556.
- 13 - Kozora E, Ellison MC, West S. Depression, fatigue, and pain in systemic lupus erythematosus (SLE): relationship to the American College of Rheumatology SLE neuropsychological battery. *Arthritis Rheum* 2006; 55:628-635.
- 14 - Govoni M, Bombardieri S, Bortoluzzi A. Factors and comorbidities associated with first neuropsychiatric event in systemic lupus erythematosus: does a risk profile exist? A large multicentre retrospective cross-sectional study on 959 Italian patients. *Rheumatology (Oxford)* 2012; 51:157-168.
- 15- Brunner HI, Ruth NM, German A. Initial validation of the Pediatric Automated Neuropsychological Assessment Metrics for childhood-onset systemic lupus erythematosus. *Arthritis Rheum* 2007; 57:1174-1182.
- 16 - Denburg SD, Denburg JA. Cognitive dysfunction and antiphospholipid antibodies in systemic lupus erythematosus. *Lupus* 2003; 12:883-890.
- 17 - Alarcón GS, Cianfrini L, Bradley LA. Systemic lupus erythematosus in three ethnic groups. X. Measuring cognitive impairment with the cognitive symptoms inventory. *Arthritis Rheum* 2002; 47:310-319.
- 18 - Ward MM, Marx AS, Barry NN. Psychological distress and changes in the activity of systemic lupus erythematosus. *Rheumatology (Oxford)* 2002; 41:184-8.
- 19 - Williams TS, Aranow C, Ross GS. Neurocognitive impairment in childhood-onset systemic lupus erythematosus: measurement issues in diagnosis. *Arthritis Care Res (Hoboken)* 2011; 63:1178-1187.

- 20 - Unterman A, Nolte JE, Boaz M, Abady M, Shoenfeld Y, Zandman-Goddard G. Neuropsychiatric syndromes in systemic lupus erythematosus: a meta-analysis. *Semin Arthritis Rheum* 2011; 41:1-11.
- 21 - Moses N, Wiggers J, Nicholas C, Cockburn J. Prevalence and correlates of perceived unmet needs of people with systemic lupus erythematosus. *Patient Educ Couns* 2005; 57:30-38.
- 22 - Iverson GL. Screening for depression in systemic lupus erythematosus with the British Columbia Major Depression Inventory. *Psychol Rep* 2002; 90:1091-1096.
- 23 - Appenzeller S, Cendes F, Costallat LT. Cognitive impairment and employment status in systemic lupus erythematosus: a prospective longitudinal study. *Arthritis Rheum* 2009; 61:680-687.
- 24 - Panopalis P, Julian L, Yazdany J. Impact of memory impairment on employment status in persons with systemic lupus erythematosus. *Arthritis Rheum* 2007; 57:1453-1460.
- 25 - Julian LJ, Yelin E, Yazdany J. Depression, medication adherence, and service utilization in systemic lupus erythematosus. *Arthritis Rheum* 2009; 61:240-246.
- 26 - Sundquist K, Li X, Hemminki K, Sundquist J. Subsequent risk of hospitalization for neuropsychiatric disorders in patients with rheumatic diseases: a nationwide study from Sweden. *Arch Gen Psychiatry* 2008; 65:501-507.
- 27 - Holliday SL, Navarrete MG, Hermosillo-Romo D. Validating a computerized neuropsychological test battery for mixed ethnic lupus patients. *Lupus* 2003; 12:697-703.
- 28 - Julian LJ, Yazdany J, Trupin L, Criswell LA, Yelin E, Katz PP. Validity of brief screening tools for cognitive impairment in RA and SLE. *Arthritis Care Res (Hoboken)* Published Online First: 12 Dec 2011. doi: 10.1002/acr.21566.

- 29 - Hanly JG, Urowitz MB, Jackson D. SF-36 summary and subscale scores are reliable outcomes of neuropsychiatric events in systemic lupus erythematosus. *Ann Rheum Dis* 2011; 70:961-967.
- 30 - Vogel A, Bhattacharya S, Larsen JL, Jacobsen S. Do subjective cognitive complaints correlate with cognitive impairment in systemic lupus erythematosus? A Danish outpatient study. *Lupus* 2011; 20:35-43.
- 31 - Hanly JG, Omisade A, Su L, Farewell V, Fisk JD. Assessment of cognitive function in systemic lupus erythematosus, rheumatoid arthritis, and multiple sclerosis by computerized neuropsychological tests. *Arthritis Rheum* 2010; 62:1478-1486.
- 32 - Adhikari T, Piatti A, Luggen M. Cognitive dysfunction in SLE: development of a screening tool. *Lupus* 2011; 20:1142-1146.
- 33 - Hawro T, Krupińska-Kun M, Rabe-Jabłońska J. Psychiatric disorders in patients with systemic lupus erythematosus: association of anxiety disorder with shorter disease duration. *Rheumatol Int* 2011; 31:1387-1391.
- 34 - dos Santos MC, Okuda EM, Ronchezel MV. Verbal ability impairment in juvenile systemic lupus erythematosus. *Rev Bras Reumatol* 2010; 50:362-374.
- 35 - Kozora E, Ellison MC, West S. Reliability and validity of the proposed American College of Rheumatology neuropsychological battery for systemic lupus erythematosus. *Arthritis Rheum* 2004; 51:810-818.
- 36 - Roebuck-Spencer TM, Yarboro C, Nowak M. Use of computerized assessment to predict neuropsychological functioning and emotional distress in patients with systemic lupus erythematosus. *Arthritis Rheum* 2006; 55:434-441.

- 37 - Kozora E, Arciniegas DB, Filley CM. Cognitive and neurologic status in patients with systemic lupus erythematosus without major neuropsychiatric syndromes. *Arthritis Rheum* 2008; 59:1639-1646.
- 38 - Zakeri Z, Shakiba M, Narouie B, Mladkova N, Ghasemi-Rad M, Khosravi A. Prevalence of depression and depressive symptoms in patients with systemic lupus erythematosus: Iranian experience. *Rheumatol Int* Published Online First: 21 Jan 2011. PubMed PMID: 21253731.
- 39 - Jarpa E, Babul M, Calderón J. Common mental disorders and psychological distress in systemic lupus erythematosus are not associated with disease activity. *Lupus* 2011; 20:58-66.
- 40 - Julian LJ, Gregorich SE, Tonner C. Using the Center for Epidemiologic Studies Depression Scale to screen for depression in systemic lupus erythematosus. *Arthritis Care Res (Hoboken)* 2011; 63:884-690.
- 41 - Nery FG, Borba EF, Viana VS. Prevalence of depressive and anxiety disorders in systemic lupus erythematosus and their association with anti-ribosomal P antibodies. *Prog Neuropsychopharmacol Biol Psychiatry* 2008; 32:695-700.
- 42 - Kozora E, Ellison MC, West S. Depression, fatigue, and pain in systemic lupus erythematosus (SLE): relationship to the American College of Rheumatology SLE neuropsychological battery. *Arthritis Rheum* 2006; 55:628-635.
- 43 - Kozora E, Arciniegas DB, Zhang L, West S. Neuropsychological patterns in systemic lupus erythematosus patients with depression. *Arthritis Res Ther* 2007; 9:R48.
- 44 - Hyphantis T, Kotsis K, Voulgari PV, Tsifetaki N, Creed F, Drosos AA. Diagnostic accuracy, internal consistency, and convergent validity of the Greek version of the

patient health questionnaire 9 in diagnosing depression in rheumatologic disorders.

Arthritis Care Res (Hoboken) 2011; 63:1313-1321.

45 - Mosca M, Govoni M, Tomietto P. The development of a simple questionnaire to screen patients with SLE for the presence of neuropsychiatric symptoms in routine clinical practice. Lupus 2011; 20(5):485-492.




Figure I Flowchart showing research process

Normas de publicação
Revista Acta Reumatológica Portuguesa

A Acta Reumatológica Portuguesa publica artigos originais sobre todos os temas da Reumatologia ou com ela relacionados. São também publicados artigos de revisão, casos clínicos, imagens, cartas ao editor e outros que se incluam na estrutura editorial da revista (recomendações, artigos sobre prática clínica reumatológica, notícias de reuniões de sociedades científicas, por ex.).

A Acta Reumatológica Portuguesa subscreve os requisitos para apresentação de artigos a revistas biomédicas elaboradas pela Comissão Internacional de Editores de Revistas Médicas (*International Committee of Medical Journal Editors*), publicada na íntegra inicialmente em N Engl J Med 1991; 324: 424-28 e actualizada em Outubro de 2008 e disponível em www.ICMJE.org. A política editorial da Acta Reumatológica Portuguesa segue as Recomendações de Política Editorial (*Editorial Policy Statements*) emitidas pelo Conselho de Editores Científicos (*Council of Science Editors*), disponíveis em www.councilscienceeditors.org/services/draft_approved.cfm.

A Revista está indexada no PubMed/Medline e os artigos estão disponíveis *online* na íntegra, com acesso aberto e gratuito.

Os artigos devem preferencialmente ser redigidos em inglês. Os artigos em língua portuguesa também podem ser submetidos para apreciação.

O rigor e a exactidão dos conteúdos, assim como as opiniões expressas são da exclusiva responsabilidade dos autores.

Os autores devem declarar potenciais conflitos de interesse.

Os artigos não podem ter sido anteriormente publicados noutra revista. Quando o artigo é aceite para publicação é mandatário o envio via e-mail de documento digitalizado, assinado por todos os autores, com a transferência dos direitos de autor para a Acta Reumatológica Portuguesa.

Os artigos publicados ficarão propriedade da revista, não podendo ser reproduzidos, no todo ou em parte, sem autorização dos editores.

A aceitação dos originais enviados para publicação é sempre condicionada a avaliação pelos consultores editoriais. Nesta avaliação os artigos poderão ser:

- aceites sem alterações;
- aceites após modificações propostas pelos revisores;
- recusados.

Em todos os casos os pareceres dos consultores serão integralmente comunicados aos autores.

Quando são propostas alterações, o autor deverá enviar via e-mail no prazo de 1 mês, uma carta ao editor e a cada um dos revisores respondendo a todas as questões colocadas e uma versão revista do artigo com as alterações inseridas destacadas com cor diferente.

Instruções aos Autores

Todos os manuscritos que não estejam em conformidade com as instruções que se seguem podem ser enviados para modificações antes de serem revistos pelos consultores.

Todos os trabalhos devem ser enviados por e-mail para: edtecnicar@gmail.com.

Os manuscritos devem ser acompanhados de declaração de originalidade e de cedência dos direitos de propriedade do artigo, assinada por todos os autores, conforme minuta publicada em anexo.

O texto deve ser enviado em formato digital (e-mail), a dois espaços, com letra tamanho 12 e com margens não inferiores a 2,5 cm, em Word para Windows. Todas as páginas devem ser numeradas.

As imagens devem ser fornecidas independentemente do texto em formato JPEG ou TIFF.

Os textos devem ser organizados da seguinte forma:

Página 1

- Título em português e inglês
- Nome dos autores e respectiva afiliação
- Serviço(s) ou organismo(s) onde o trabalho foi executado
- Subsídio(s) ou bolsa(s) que contribuíram para a realização do trabalho
- Morada e e-mail do autor responsável pela correspondência relativa ao manuscrito
- Título breve para rodapé

Página 2

- Título (sem autores)
- Resumo em português e inglês, que para os artigos originais deve ser estruturado da seguinte forma: Objectivos, Material e Métodos, Resultados, Conclusões. O resumo dos artigos originais não deve exceder as 350 palavras e o dos casos clínicos as 180 palavras.
- Palavras-chave em português e em inglês (Keywords)

Um máximo de 5 palavras-chave, utilizando a terminologia que consta na lista do Index Medicus: «Medical Subject Headings» (MeSH), deve seguir-se ao resumo.

Página 3 e seguintes

Artigos originais: O texto deve ser apresentado com os seguintes subtítulos: Introdução (incluindo Objectivos), Material e Métodos, Resultados, Discussão, Conclusões, Agradecimentos (se aplicável), Referências.

Os artigos originais não deverão exceder as 4.000 palavras, com um total de 6 figuras/tabelas e 60 referências.

Caso clínico: os subtítulos serão, Introdução, Caso clínico, Discussão, Referências.

O caso clínico não deve exceder as 2.000 palavras e 25 referências. Deve ser acompanhado de figuras ilustrativas. O número de tabelas/figuras não deve ser superior a 6.

A partir da segunda página, inclusive, todas as páginas devem ter em rodapé o título breve indicado na página 1.

Referências: As referências bibliográficas devem ser classificadas e numeradas por ordem de entrada no texto, em *superscript* e não entre parêntesis. As abreviaturas usadas na nomeação das revistas devem ser as utilizadas pelo *Index Medicus*.

Nas referências com 6 ou menos autores todos devem ser nomeados. Nas referências com 7 ou mais autores devem ser nomeados os 3 primeiros seguidos de et al.

Notas: Os números da página inicial e final devem ser totalmente apresentados (565-569 e não 565-9)

Não indicar o número da revista nem o mês da publicação.

Seguem-se alguns exemplos de como devem constar os vários tipos de referências:

– *Revista*

Apelido e iniciais do(s) autor(es). Título do artigo. Nome da revista Ano; Volume: Páginas.

Ex.: Hill J, Bird HA, Hopkins R, Lawton C, Wright V. Survey of satisfaction with care in a rheumatology outpatient clinic. *Ann Rheum Dis* 1992; 51:195-197.

– *Artigo publicado online (inserir DOI)*

Ex.: Peter A Merkel, David Curthbertson, Bernhard Hellmich et al. Comparison of disease activity measures for ANCA-associated vasculitis. *Ann Rheum Dis* Published Online First: 29 July 2008. doi:10.1136/ard.2008.097758

– *Capítulo de livro*

Nome(s) e iniciais do(s) autor(es) do capítulo. Título do capítulo. In: Nome(s) e iniciais do(s) editor(es) médico(s). Título do livro. Cidade: Nome da casa editora, ano de publicação: primeira a última página do capítulo.

Ex.: Stewart AF. Hypercalcemia resulting from medications. In: Favus MJ, ed. *Primer on the Metabolic Bone Diseases and Disorder of Mineral Metabolism*. New York: Raven Press, 1993: 177-178.

– *Livro*

Nome(s) e iniciais do(s) autor(es). Título do livro. Cidade: Nome da casa editora, ano de publicação: página(s).

Ex.: Lorig K. *Patient Education. A practical approach*. St. Louis: Mosby-Year Book;1992: 51.

– *Documento electrónico*

Ex: Programa Nacional de Luta Contra a Tuberculose. Sistema de Vigilância (SVIG-TB). Direcção-Geral da Saúde - Divisão de Doenças Transmissíveis, Março de 2005 <http://www.dgsaude.pt/upload/membro.id/ficheiros/i006875.pdf>. Acedido em 25 Janeiro de 2008

As referências a trabalhos ainda não publicados, comunicações em reuniões, não publicadas em livros de resumos, ou comunicações pessoais devem ser citadas no texto e não como referências formais.

A exactidão e o rigor das referências são da responsabilidade do autor.

Tabelas: As tabelas a inserir devem ser assinaladas no texto em numeração romana e cumprir o limite descrito acima. Cada tabela deverá ser apresentada em folha separada, dactilografada a 2 espaços. Na parte superior devem apresentar um título sucinto mas informativo, de modo a poder ser compreendido sem recurso ao texto. Na parte inferior da tabela deve constar a explicação das abreviaturas utilizadas. Nas tabelas devem ser evitados os traços verticais e os traços horizontais, estes devem servir apenas como separadores de títulos e subtítulos.

Figuras: As figuras a inserir devem ser assinaladas no texto em numeração árabe e cumprir o limite definido acima. As legendas das figuras devem ser dactilografadas a dois espaços numa folha separada, depois da bibliografia. As figuras devem ser enviadas em suporte informático com ficheiros separados para cada figura, em formato JPEG ou TIFF.

Editoriais: Os editoriais serão solicitados por convite do Editor. Os editoriais serão comentários sobre tópicos actuais ou sobre artigos publicados na revista. O texto dos

editoriais não deve exceder as 1.200 palavras, um máximo de 15 referências e não deve conter quadros ou figuras.

Artigos de revisão: Estes artigos serão preferencialmente solicitados pelo Editor. No entanto, os autores interessados em apresentar um artigo de revisão podem contactar o Editor para discussão dos tópicos a apresentar.

O artigo de revisão não deve exceder as 4.000 palavras, 6 tabelas/figuras e 100 referências.

Cartas ao Editor: As cartas ao editor devem constituir um comentário crítico a um artigo da revista ou uma pequena nota sobre um tema ou caso clínico. Não devem exceder as 600 palavras, uma figura ou um quadro, e um máximo de 10 referências bibliográficas.

Imagens em reumatologia: Podem ser submetidas imagens de particular interesse. As figuras, no máximo de 4, devem ser enviadas em formato JPEG ou TIFF de boa resolução. O texto acompanhante não deve ultrapassar as 500 palavras.

Modificações e revisões: No caso da aceitação do artigo ser condicionada a modificações, estas deverão ser feitas pelos autores no prazo de 1 mês.

Quando são propostas alterações, o autor deverá endereçar uma carta ao editor e a cada um dos revisores respondendo a todas as questões colocadas. Deverá ainda submeter uma versão revista do artigo com as alterações inseridas destacadas com cor diferente.

As provas tipográficas serão, sempre que possível, enviadas aos autores contendo a indicação do prazo para revisão consoante as necessidades editoriais da revista.

Minuta da carta de submissão a enviar ao Editor, digitalizada, por e-mail:

Enviar este documento com o manuscrito para:
edtecnicarp@gmail.com

Editor

Acta Reumatológica Portuguesa

O(s) autor(es) certifica(m) que o manuscrito intitulado:

(ref. ARP _____) é original, que todas as afirmações apresentadas como factos são baseados na investigação do(s) autor(es), que o manuscrito, quer em parte quer no todo, não infringe nenhum *copyright* e não viola nenhum direito da privacidade, que não foi publicado em parte ou no todo e que não foi submetido para publicação, no todo ou em parte, noutra revista, e que os autores têm o direito ao *copyright*.

Todos os autores declaram ainda que participaram no trabalho, se responsabilizam por ele e que não existe, da parte de qualquer dos autores conflito de interesses nas afirmações proferidas no trabalho.

Os autores, ao submeterem o trabalho para publicação, transferem para a Acta Reumatológica Portuguesa todos os direitos a interesses do *copyright* do artigo.

Todos os autores devem assinar

Data: _____

Nome (maiúsculas): _____

Assinatura: _____